



Physical and Artificial Resistivity (in smoothed particle magnetohydrodynamics)

James Wurster


1st Phantom Users Workshop
Monash University, 20 February 2018




Ideal magnetohydrodynamics





$$\frac{d\mathbf{B}}{dt} = (\mathbf{B} \cdot \nabla) \mathbf{v} - \mathbf{B} (\nabla \cdot \mathbf{v})$$




Ideal MHD

- Fully ionised plasma 
- Zero resistivity & infinite conductivity
- Ions & electrons are tied to the magnetic field







Ideal MHD



Density (rendered) + Magnetic field lines
Ideal MHD. Left: Initial conditions. Right: at $\rho_{\max} = 10^{-9} \text{g cm}^{-3}$




Ideal MHD: Artificial Resistivity



$$\begin{aligned}\frac{d\mathbf{B}}{dt} &= (\mathbf{B} \cdot \nabla) \mathbf{v} - \mathbf{B} (\nabla \cdot \mathbf{v}) \\ &+ \nabla \times \eta_{\text{art}} (\nabla \times \mathbf{B})\end{aligned}$$

where

$$\eta_{\text{art}} \approx \frac{1}{2} \alpha_B v_{\text{sig}} h$$



Ideal MHD: Artificial Resistivity

➤ Artificial resistivity (Tricco & Price, 2013)

$$\frac{dB_a^i}{dt} = -\frac{1}{\Omega_a \rho_a} \sum_b m_b \left[v_{ab}^i B_a^j \nabla_a^j W_{ab}(h_a) - B_a^i v_{ab}^j \nabla_a^j W_{ab}(h_a) \right] + \left. \frac{dB_a^i}{dt} \right|_{\text{art}}$$

$$\left. \frac{dB_a^i}{dt} \right|_{\text{art}} = \frac{\rho_a}{2} \sum_b m_b B_{ab}^i \left[\frac{\alpha_a^B v_{\text{sig},a} \hat{r}_{ab}^j \nabla_a^j W_{ab}(h_a)}{\Omega_a \rho_a^2} + \frac{\alpha_b^B v_{\text{sig},b} \hat{r}_{ab}^j \nabla_a^j W_{ab}(h_b)}{\Omega_b \rho_b^2} \right]$$

$$v_{ab}^i = v_a^i - v_b^i$$


$$B_{ab}^i = B_a^i - B_b^i$$

$$v_{\text{sig},a} = \sqrt{c_{s,a}^2 + v_{A,a}^2}$$

$$\alpha_a^B = \min \left(\frac{h_a |\nabla \mathbf{B}_a|}{|\mathbf{B}_a|}, 1 \right)$$

$$|\nabla \mathbf{B}_a| \equiv \sqrt{\sum_i \sum_j \left| \frac{\partial B_a^i}{\partial x_a^j} \right|^2}$$

➤ Always applied if there is a gradient in the magnetic field (i.e. $|\nabla \mathbf{B}| > 0$)



Ideal MHD: Artificial Resistivity

- Artificial resistivity (Price, et al, submitted)

$$\frac{dB_a^i}{dt} = -\frac{1}{\Omega_a \rho_a} \sum_b m_b \left[v_{ab}^i B_a^j \nabla_a^j W_{ab}(h_a) - B_a^i v_{ab}^j \nabla_a^j W_{ab}(h_a) \right] + \left. \frac{dB_a^i}{dt} \right|_{\text{art}}$$


$$\left. \frac{dB_a^i}{dt} \right|_{\text{art}} = \frac{\rho_a}{2} \sum_b m_b \alpha^B v_{\text{sig},ab} B_{ab}^i \left[\frac{\hat{r}_{ab}^j \nabla_a^j W_{ab}(h_a)}{\Omega_a \rho_a^2} + \frac{\hat{r}_{ab}^j \nabla_a^j W_{ab}(h_b)}{\Omega_b \rho_b^2} \right]$$

$$B_{ab}^i = B_a^i - B_b^i$$

$$v_{\text{sig},ab} = |\mathbf{v}_{ab} \times \hat{\mathbf{r}}_{ab}|$$

$$\alpha^B \equiv 1$$

- Always applied for non-zero velocity
- Less resistive than that from Tricco & Price (2013)



Ideal MHD: Artificial Resistivity

- Price et. al. (2017) artificial resistivity

$$v_{\text{sig},ab} = |\mathbf{v}_{ab} \times \hat{\mathbf{r}}_{ab}|$$


$$\alpha^{\text{B}} \equiv 1$$

- Tricco & Price (2013)



$$v_{\text{sig},a} = \sqrt{c_{s,a}^2 + v_{A,a}^2}$$


$$\alpha_a^{\text{B}} = \min \left(\frac{h_a |\nabla \mathbf{B}_a|}{|\mathbf{B}_a|}, 1 \right)$$

- Tricco & Price (2013) with alternate averaging



Ideal MHD: Artificial Resistivity








Motivation: Non-ideal MHD




Orion Molecular Cloud
Ionisation fraction $\sim 10^{-14}$






HL Tau
 $\sim 10^{-12}$



Non-ideal MHD

- Partially ionised plasma 
- Non-zero resistivity & conductivity
- Ions, electrons & neutrals behaviour is environment-dependent







$$\left. \frac{dB}{dt} \right|_{\text{OR}} = -\nabla \times \eta_{\text{OR}} (\nabla \times B),$$


$$\left. \frac{dB}{dt} \right|_{\text{HE}} = -\nabla \times \eta_{\text{HE}} [(\nabla \times B) \times \hat{B}],$$

$$\left. \frac{dB}{dt} \right|_{\text{AD}} = \nabla \times \eta_{\text{AD}} \left\{ [(\nabla \times B) \times \hat{B}] \times \hat{B} \right\}.$$




Non-ideal MHD: Hall effect







Ideal vs non-ideal MHD





Density (rendered) + Magnetic field lines
During first core phase. Left: ideal MHD. Right: non-ideal MHD



Ideal vs non-ideal MHD



Density (rendered) + Magnetic field lines
During first core phase. Left: ideal MHD. Right: non-ideal MHD



$$\left. \frac{dB}{dt} \right|_{\text{OR}} = -\nabla \times \eta_{\text{OR}} (\nabla \times B),$$

$$\left. \frac{dB}{dt} \right|_{\text{HE}} = -\nabla \times \eta_{\text{HE}} [(\nabla \times B) \times \hat{B}],$$

$$\left. \frac{dB}{dt} \right|_{\text{AD}} = \nabla \times \eta_{\text{AD}} \left\{ [(\nabla \times B) \times \hat{B}] \times \hat{B} \right\}.$$

NICIL: Wurster (2016)

Marchand+ (2016)

NICIL v1.2.3 is implemented in the current git version of Phantom

Non-ideal MHD in Phantom: the NICIL library

- *Phantom* includes the *NICIL* code (Wurster 2016)
 - Publically available at <https://bitbucket.org/jameswurster/nicil>
- When compiling, set `NONIDEALMHD=yes`
- Realistic defaults are set; these will self-consistently calculate the non-ideal coefficients
- Fully parameterisable
- Primary parameters are included in *Phantom*'s `.in` file
- All parameters are included at the top of `nicil.F90`
- Important parameters that can be modified
 - Included non-ideal MHD terms (default = ohmic + Hall + ambipolar)
 - Ionisation source (default = cosmic rays + thermal)
 - Cosmic ray ionisation rate (default = 10^{-17} s^{-1})
 - Elements that can be thermally ionised (cannot be modified through `.in` file)
 - Grain properties (default = fixed size of $0.1 \mu\text{m}$; alternate is MRN, but is slow)
- Important values are summarised in the dump files and the `.ev` file
- Can optionally preselect non-ideal MHD coefficients (preferably for tests only)
- All coefficients and required variables are calculated at runtime

Implementation

➤ Continuum equations

$$\begin{aligned}
 \frac{d\mathbf{B}}{dt} = & (\mathbf{B} \cdot \nabla) \mathbf{v} - \mathbf{B} (\nabla \cdot \mathbf{v}) \\
 & + \nabla \times \eta_{\text{art}} (\nabla \times \mathbf{B}) \\
 & + \nabla \times \eta_{\text{OR}} (\nabla \times \mathbf{B}) \\
 & + \nabla \times \eta_{\text{HE}} \left[(\nabla \times \mathbf{B}) \times \hat{\mathbf{B}} \right] \\
 & + \nabla \times \eta_{\text{AD}} \left\{ \left[(\nabla \times \mathbf{B}) \times \hat{\mathbf{B}} \right] \times \hat{\mathbf{B}} \right\}
 \end{aligned}$$

➤ SPMHD equations

$$\frac{dB_a^i}{dt} = -\frac{1}{\Omega_a \rho_a} \sum_b m_b \left[v_{ab}^i B_a^j \nabla_a^j W_{ab}(h_a) - B_a^i v_{ab}^j \nabla_a^j W_{ab}(h_a) \right] + \left. \frac{dB_a^i}{dt} \right|_{\text{non-ideal}}$$


$$\left. \frac{d\mathbf{B}_a}{dt} \right|_{\text{non-ideal}} = -\rho_a \sum_b m_b \left[\frac{\mathbf{D}_a}{\Omega_a \rho_a^2} \times \nabla_a W_{ab}(h_a) + \frac{\mathbf{D}_b}{\Omega_b \rho_b^2} \times \nabla_a W_{ab}(h_b) \right],$$

$$\mathbf{D}_a^{\text{OR}} = -\eta_{\text{OR}} \mathbf{J}_a, \quad \mathbf{D}_a^{\text{HE}} = -\eta_{\text{HE}} \mathbf{J}_a \times \hat{\mathbf{B}}_a, \quad \mathbf{D}_a^{\text{AD}} = \eta_{\text{AD}} (\mathbf{J}_a \times \hat{\mathbf{B}}_a) \times \hat{\mathbf{B}}_a.$$

Implementation


Density Loop:

```
do i = 1, N
  do j = 1, N_neigh
    Using j, calculate density of i
    Using j, calculate current density,  $\mathbf{J} = \nabla \times \mathbf{B}$ , of i
  enddo
  Using new density of i, calculate  $\eta_{\text{nimhd}}$ 
enddo
```



Force Loop:

```
do i = 1, N
  Calculate  $\mathbf{J}_i \times \mathbf{B}_i$  and  $(\mathbf{J}_i \times \mathbf{B}_i) \times \mathbf{B}_j$ 
  do j = 1, N_neigh
    Calculate  $\mathbf{J}_j \times \mathbf{B}_j$  and  $(\mathbf{J}_j \times \mathbf{B}_j) \times \mathbf{B}_i$ 
    Using j, calculate  $d\mathbf{B}/dt_{\text{non-ideal}}$  of i
  enddo
  Calculate non-ideal timesteps
enddo
```



Step Loop:

```
do i = 1, N
  Updated magnetic field of i, using ideal, non-ideal and artificial terms
enddo
```

Implementation

➤ Timestepping:


$$dt_{\text{Courant}} = C_c \frac{h}{v_{\text{sig}}}$$

$$dt_{\text{nimhd}} = C_{\text{ni}} \frac{h^2}{|\eta|}$$

➤ *Phantom* includes super-timestepping (Alexiades, Amiez & Gremaud 1996)

➤ Right: cpu-hours required for the 10^6 particle models with $\mu_0=5$ in Wurster, Price & Bate (2016)

➤ Non-ideal MHD is slightly slower for $t < t_{\text{ff}}$, and much slower for $t > t_{\text{ff}}$





Conclusions


- Artificial resistivity is required to stabilised magnetohydrodynamics equations
- Ideal MHD is a poor approximation for modelling molecular clouds or protoplanetary discs
- Non-ideal MHD requires an assumption of chemistry
- The non-ideal MHD coefficients are not dependent on neighbours
- The non-ideal MHD contribution to the magnetic field evolution is dependent on neighbours
- Non-ideal MHD introduces a diffusion timestep $\propto h^2$, hence can be computationally expensive

```
return 'N/A'
```

"Always code as if the guy who ends up
maintaining your code will be a violent
psychopath who knows where you live."

```
suffix = ['B', 'KiB', 'MiB', 'GiB', 'TiB', 'PiB', 'EiB', 'ZiB', 'YiB'] * [exponent]  
converted = float(bytes) / float(1024 ** exponent)  
return '%.2f%s' % (converted, suffix)
```

~ John Woods



j.wurster@exeter.ac.uk

<http://www.astro.ex.ac.uk/people/wurster/>

Presentation available at http://www.astro.ex.ac.uk/people/wurster/files/spmhd_resistivity.pdf

Nicil's git repository: <https://bitbucket.org/jameswurster/nicil>